

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

BROWN et al

Atty. Ref.: 4598-2; Confirmation No. 2642

Appl. No. 10/723,420

TC/A.U. 1614

Filed: November 26, 2003

Examiner: Royds, L.

For: BIOLOGICALLY ACTIVE METHYLENE BLUE DERIVATIVES

* * * * *

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

DECLARATION

I, DAVID M. LEWIS, do hereby declare and state as follows:

1. I have a PhD degree in Colour Chemistry and I have held the position of Professor and Head of the Department from 1987 to 2004, and Emeritus Professor from 2004 to the present time, in the Department of Colour Chemistry at the University of Leeds. In my 40 years experience in industry and academia as a dye chemist, I have gained an in-depth knowledge of the chemistry of dyes and their applications. My research experience includes studies of the relationships between dye structure and dye properties, the mechanistic and technological aspects of dye uptake by natural and synthetic polymers, dye photochemistry, modified polymers and bactericidal agents. A copy of my curriculum vitae is attached.
2. I have reviewed the contents of U.S. Application No. 10/723,420 and the claims presently on file, specifically, claims 77-79, 84, 89-91, and 98 - 100. I have also reviewed the

Office Action dated 25th June 2007, and understand that the Examiner has rejected the claims on the basis that they would have been obvious over various prior art references [Note 1 (b) - (h)]. The Examiner suggests that the prior art references which are directed to the use of Methylene Blue which is of identical core structure to the presently claimed compounds would have rendered the claims obvious because the skilled person would have reasonably predicted that the present compounds would share similar pharmacologic properties with those of Methylene Blue due to their structural homology. Further, the Examiner suggests that the skilled person would have expected the present compounds to have exhibited bactericidal activity against the claimed bacteria the same or substantially similar to that exhibited by Methylene Blue.

3. The cationic dye Methylene Blue has been known since 1876, and since the end of the 19th century has been investigated as an antimicrobial agent for several therapeutic applications [see for example **Ref. 1**, p.99]. It is notable that the photoantimicrobial activity of Methylene Blue is reported as early as 1930 [**Ref. 2** p.17], and yet despite the obvious rewards of producing improved analogues. I am not aware of any report of any higher alkyl homologues (ethyl, propyl, butyl etc.), or homologues with other hydrophobic side chains, ever having been investigated as antimicrobial agents prior to the filing of U.S. Application No. 10/723,420. This suggests that there were good scientific reasons why researchers were not led to investigate these higher hydrophobic homologues prior to the present invention. M. Wainwright in his review of photodynamic therapy (**Ref 1**, p. 100) states: "*Phenothiazinium dyes, usually based on the Methylene Blue structure, are eminently suited to use as biomedical photosensitisers. ... Unfortunately the optimal nature of the lead compound (i.e. Methylene Blue) has often discouraged the search for improved analogues.*"

It is my opinion that conventional scientific thinking on the mechanism of cationic dye staining of polymers (i.e. microorganisms), which is one of attraction between oppositely charged ions [see for example **Ref. 3**, pp 2-3], has in the past led the skilled person away from using more hydrophobic higher homologues of Methylene Blue as photo-antibacterial agents of equivalent activity to Methylene Blue. In my opinion, existing knowledge would have made it unobvious that such Methylene Blue homologues would have had any advantages over Methylene Blue itself, and indeed the literature would have suggested that there would be a greater likelihood of such homologues having therapeutic properties inferior to those of Methylene Blue. My reasons for this opinion are as follows:

It was generally accepted that photo-antimicrobial cationic dyes such as Methylene Blue worked by first staining biological components of the cell, and then, after exposure to light, generating singlet oxygen which attacked cell constituents and caused rapid cell death. [See for example M. Wainwright, **Ref. 4**, p. 351 “*If a known synthetic photosensitizer such as toluidine blue could be seen to stain a pathogenic organism sufficiently, then it should be possible to destroy that organism via subsequent irradiation.*”; M. Wainwright, **Ref. 2**, p 14. “*It is this work which underlies the principle of photodynamic antimicrobial action—if a live microbe could be demonstrated selectively with a vital stain such as methylene blue, it should be possible to destroy the stained microbe on illumination*”]. Thus, efficient adsorption of the dye onto the cell components in contact with the dye solution, i.e. cell staining, was seen to be one of the most important characteristics of a therapeutically effective photo-antimicrobial agent.

This selective *cell staining* by cationic dyes (such as Methylene Blue) has been used for many years [see for example **Ref. 4**, page 355] in the histological study of microorganisms. *Cationic dyeing of textiles* operates by an identical process to cell staining, and similarly involves selective uptake of a cationic dye from solution by an anionic polymer, such as protein fibers (e.g. wool, silk) and polyacrylic fibres modified with anionic groups. Thus, the effectiveness of a cationic dye as a photo-antimicrobial agent was understood to be related to its effectiveness both as a biological stain and as a dye for anionic fibers.

It was also understood [see for example **Ref. 5**] that cell staining with cationic dyes was most successfully achieved using an aqueous solution of the dye. Almost all microbiological stain solutions based on ionic dyes use water as the preferred solvent. Ethanol is sometimes used, but gives poorer uptake [**Ref. 5**]. Water was viewed as the ideal solvent for delivery of a cationic photo-antimicrobial dye to microorganisms in a therapeutic situation for the following reasons:

- (a) Uptake of a cationic dye at the negatively charged cell surface involves ion exchange, in which cations on the biopolymer (H^+ Na^+ etc.) are replaced by the dye cation. The dye cation then becomes paired with an anionic group on the surface. [**Ref. 3**, pages 2-3]. Water is a unique solvent, and its exceptionally high dielectric constant and solvating power ensures dissociation of ion pairs. The ion exchange process thus proceeds with much greater facility in water than in organic solvents.
- (b) Alternative organic solvents (usually ethanol) are often unsuitable for staining with cationic dyes, and give poorer uptake when the staining mechanism is ionic in nature [**Ref. 5**].

(c) An additional important factor which made water the solvent of choice for photo-antimicrobial dyes is that it is the only solvent that could be used safely in undiluted form for therapeutic application to living tissue. Organic solvents (such as ethanol) could only be used if diluted with water to low concentrations because of their inherent adverse effects on the host tissue.

Thus the skilled person developing cationic dye photosensitizers for therapy would have been led to water as the ideal vehicle for delivering these dyes to the target microorganism.

In staining or dyeing, if a cationic dye is to be adsorbed from an aqueous solution onto a cell or textile fiber surface, then good water solubility had long been regarded as an important prerequisite, as the rate of adsorption was expected to be dependent on the concentration of the dye in the aqueous phase. Thus, it had long been accepted that cationic dyes for textile dyeing needed to have good water solubility so that solution concentrations conducive to rapid dye uptake could be used. Dyes that had low water solubility as a consequence of hydrophobic side chains were expected to produce only weak aqueous solutions when fully saturated and such weak solutions were expected to result in unacceptably slow uptake rates. This perceived good water solubility requirement is clearly indicated by an examination of the **Colour Index [Ref. 6]** which lists all known dyes that have ever been commercialized. Consideration of the 3rd Edition (1971) shows that more than 150 synthetic cationic dyes of known structure had been commercialized up to that time, and all of these had the common characteristic of good water solubility. Representative examples from different chemical classes are: the oxazine Basic Blue 3 (solubility 3g/100ml), the rhodamine Basic Red 1 (solubility 2 g/100ml), the thiazole Basic

Yellow 1 (solubility 2g/100ml; the triphenylmethane Basic Violet 1 (solubility 5g/100ml), and Methylene Blue itself (Basic Blue 9) (solubility 5 g /100ml). [Solubility data taken from **Ref. 7**]. All N-alkylated cationic dyes listed by the **Colour Index** had alkyl groups no longer than two carbon atoms and most had methyl groups, clearly suggesting the need for hydrophobic side chains to be kept to a level where water solubility was not compromised. The only exception is the azine dye CI 11090 which has a dihydroxypropyl side chain, and in this case the two adjacent hydrophilic hydroxy groups are present to counteract the insolubilising effect of the longer propyl chain. Focusing attention on the phenothiazinium class, the **Colour Index** shows that just nine Methylene Blue type structures have been commercialized since 1876 to the present day. All these compounds are anticipated to be very water soluble, as none have alkyl chains longer than ethyl. Water solubility data could be found for six of these [**Ref. 7**], and as expected, values are high and fall in the range 1-5 g/100 ml.

By analogy, cationic dyes for microbial staining would have been expected to have good water solubility, and this is also confirmed by the commercial record. The Sigma Aldrich catalogue [**Ref.8**] lists the widest range of microbiological stains in the world, many certified by the Biological Stains Commission, and includes six cationic phenothiazinium dyes among the 300 or so stains listed. These six (Methylene Blue, Azure A, Azure B, Azure C, Toluidine Blue and New Methylene Blue) are also listed in the **Colour Index**, and as noted had water solubilities in the range 1-5 g/100ml.

The marked absence of longer chain hydrophobic analogues of Methylene Blue from the historical commercial textile dye record and from microbiological stain listings could not have been due to any lack of availability of such compounds, since synthetic routes to such homologues had been readily available for more than 130 years (see for examples references 9

and 10). The absence of these compounds would have been due to their actual or perceived inferior dyeing and staining properties compared to Methylene Blue itself, such inferiority having been expected to stem from their lower water solubility.

The pronounced effect of increasing hydrophobicity on water solubility in the Methylene Blue series is exemplified by the following data for alkyl derivatives:

Solubilities in water at ca. 25 °C:

<i>Methylene Blue</i>	<i>~ 5 g / 100ml</i>
<i>Ethylene Blue</i>	<i>~ 3.9 g / 100ml</i>
<i>Propylene Blue</i>	<i>~ 0.01 g / 100ml</i>
<i>Butylene Blue</i>	<i>~ 0.015 g / 100ml</i>
<i>Pentylene Blue</i>	<i>~ 0.0001 g / 100ml</i>

[Data for Methylene Blue from Ref. 7. Data for other dyes measured by Photopharmica]

Thus, the propyl and butyl derivatives show an approximately 500 fold decrease in solubility compared to Methylene Blue, and the pentyl derivative a four orders of magnitude reduction. Such low water solubilities would have been expected to preclude these compounds from having any technical value as textile dyes, and would have suggested that they would had little value as aqueous biological stains. It was therefore most surprising that these compounds had such dramatically greater photo-antimicrobial activities than Methylene Blue and Ethylene Blue.

Thus, the relative cell kill efficiencies against *E. coli* are:

Methylene blue	3.5
Ethylene Blue	1.7
Propylene Blue	8.1
Butylene Blue	52,000
Pentylene Blue	195,000

This major enhancement in photo-antibacterial properties is even more surprising when one takes into account that there is a small decrease in activity when the homologous series is ascended from Methylene Blue to Ethylene Blue.

In conclusion, it is my opinion that long-established conventional thinking on the relationship between the water solubility of a cationic dye and its effectiveness both as a textile dye and as a biological stain would have led the skilled person away from considering more hydrophobic higher homologues of Methylene Blue, such as those claimed in US Appln. No.10/723420 as useful photo-antimicrobial dyes. Convention would have taught that such an increase in molecular size, where the increase in hydrophobicity resulted in a significant decrease in water solubility of the dye, would be detrimental to dye uptake and thus detrimental to the observed photo-antimicrobial activity. On this basis the skilled person would not have been motivated to consider higher homologues. Additionally, even if an artisan were to have considered higher homologues he/she would have expected poorer results rather than the significantly improved results obtained in the present invention. This is particularly the case if one considered the activities of Methylene Blue and Ethylene Blue. A researcher systematically investigating the effect of increased alkyl chain length and increasing hydrophobicity on activity would have found that extending the chain length from methyl to ethyl resulted in a decrease in activity, and so would have been dissuaded from examining longer chains.

There has been a long felt medical need for powerful antimicrobial drugs which operate by a mechanism different from that of antibiotics: drugs that could be used topically on

infected wounds and ulcers, had broad spectrum activity even against "superbugs", and are unlikely to lead to drug-resistant strains of microorganism. Despite the fact that photo-antibacterial therapy has long been considered a way of achieving these objectives, none of the extensive research that has taken place prior to U.S. Application No. 10/723,420 succeeded in producing new photosensitizers with the high photo-antimicrobial activities of the present compounds.

* * * * *

I hereby declare that all statements made herein of my own knowledge are true and that statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Further, declarant sayeth not.

Signed this 16th day of August, 2007.

David M. Lewis
David M. Lewis

NOTE 1:

Prior art cited in Office Action dated 25th June 2007:

- (a) Y. Mazur et al. "*Phenothiazinium salts and their use for disinfecting aqueous effluents*".
U.S. Patent No. 5,220,009; (1993).
- (b) S J Wagner "*Decontamination of whole blood and cellular components by phenothiazin-5-ium dyes plus light*". WO/91/16911 (1991)
- (c) M Wainwright et al "*Photobactericidal activity of phenothiazinium dyes against methicillin-resistant strains of Staphylococcus aureus*". FEMS Microbiology Letters, vol.160, No. 2 (1998) pp 171-181.
- (d) E. Shanbrom "*Microbicide treated polymeric materials.*" U.S. Patent No.6,183,764;(2001).
- (e) M. Wilson and W. Harvey, "*Laser treatment*". U.S. Patent No. 5,611,793 (1997)
- (f) M A Biel . "*Photodynamic cellular and acellular organism eradication utilizing a photosensitizing material and surfactant*". WO 01/62289 (2001).
- (g) R K Chowdhary and D Dolphin. "*Supports for photosensitizer formulations*" U.S.Patent Application Publication 2002/0061330 (2002).
- (h) *The Merck Index*, Monograph 5979 (1989).

REFERENCES

1. M Wainwright, *Review of Progress in Colouration*, (2004), **34** pp. 95-109].
2. M Wainwright, "Photodynamic antimicrobial chemotherapy (PACT)", *Journal of Antimicrobial Chemotherapy* (1998) **42**, 13-28
3. J.A. Kiernan, "Dyes and other colorants in microtechnique and biomedical research" *Coloration Technology*, (2006) **122**, 1-21.
4. Mark Wainwright "Non-porphyrin Photosensitizers in Biomedicine" *Chemical Society Reviews*, 1996, 351-359 .
- ✓ 5. **Stains File**, internet resource for histotechnologists,
<http://stainsfile.info/StainsFile/theory/solvent.htm>).
6. **The Colour Index 3rd. Edition 1971.**(Society of Dyers and Colorists / Americould Association of Textile Chemists and Colorists)
Current on-line version: <http://www.colour-index.org>
7. *The Sigma-Aldrich Handbook of Stains, Dyes and Indicators*, F.J. Green (Aldrich Chemical Company Inc., Milwaukee, 1990)
8. <http://www.sigmaaldrich.com>
[http://www.sigmaaldrich.com/Area_of_Interest/Research_Essentials/Hematology_and_Histology/Product_Lines/Stains_and_Dyes.html.]
9. A. Bernthsen, „Studien in der Methylenblaugruppe“ *Justus Liebig's Annalen Der Chemie*, (1888) **251**, Abschnitt 6, „Homologe Indamine und Thioninfarbstoffe“, pp 83-97.
10. J C V P Moura and N Cordeiro "Synthetic routes to 3,7-bis(dialkylamino)-phenothiazin-5-ium compounds." *Current Drug Targets*; (2003), **4** (2), pp 133-142

CURRICULUM VITAE

Professor David M. Lewis, BSc, PhD, FRCS

Business Address: Department of Colour Chemistry
University of Leeds
Leeds, LS2 9JT
UK

Telephone: (+44) (0)113 343 2931

Fax: (+44) (0)113 343 2947

Research experience: 40 years experience in industry and academia, in the general areas of dye chemistry and dye application technology. 215 refereed publications in the fields of dye chemistry, dyeing technology, dye photochemistry, polymer chemistry, analytical chemistry and ink technology. 38 patents in related fields.

Current positions:

- Emeritus Professor, Department of Colour Chemistry, The University of Leeds, Leeds LS25 9JT, UK.
- Research Director, Inovink Ltd.
- Chief Scientific Officer, Perachem Ltd.

Previous employment history:

1966 – 1978: Development Officer - Principal Development Officer, International Wool Secretariat (IWS) Ilkley. Development of new dyeing and chemical finishing processes for wool and the part supervision of chemical aspects of the Wool Foundation Northern Hemisphere sponsored research scheme.

Jan 1978 – April 1979: Senior Research Scientist, CSIRO, Division of Textile Industry, Geelong, Australia. Development of metallisable, sublimable disperse dyes for the transfer printing of wool and wool-blend fabrics.

April 1979 – April 1987: Principal Development Scientist, IWS. Coloration and chemical finishing developments..

April 1987-August 2003. Professor and Head of Department of Colour Chemistry, University of Leeds.

August 1998 – August 2003: Head of Resource Centre, School of Physical Sciences, University of Leeds.

Honours and Achievements:

Silver medal of the Society of Dyers and Colourists for Technological Achievement, March 1985.

Chairman of the West Riding Region, Society of Dyers and Colourists, March 1986 – March 1988.

Fellowship of the Royal Society of Chemistry, awarded in 1984.

Visiting Professorships in China at the North West Textile Institute (Xi'an), the University of Heilongjiang (Harbin) and the Technical University of Wuhan.

Chairman, International Association of Textile Chemists and Colourists Studentship Committee from June 1989.

Gold Research Medal, Worshipful Company of Dyers: for papers published on Reactive Dyes (January 1992).

President, Society of Dyers and Colourists, April 1993 – April 1994.

Gold Medal of the Worshipful Company of Dyers, April 2005.

Millson Award for Invention, by the American Association of Textile Chemists and Colourists, 2005.

Accreditation/Validation Experience:

BSc (Hons) Course Validator at Scottish College of Textiles, April 1987.

External Examiner, BSc (Hons) Applied Chemistry, Scottish College of Textiles, April 1987 – September 1990.

BSc (Hons) Course Validator at Hong Kong Polytechnic, April 1991.

External Examiner, BSc (Hons) Textile Chemistry, Hong Kong Polytechnic (1991/92 – 1994/95).

Science assessor for CSIRO review group, Geelong, Australia – March 2007.

LIST OF PUBLICATIONS

1. Reactive Dyes and Resin Treated Wool - Keys to the Attainment of Superwash Standards, Cockett, K R F and Lewis, D M, *J.S.D.C.*, **92** (1976) 141
2. The Use of Bunte Salt Terminated Polymers in Wool Finishing with Special Reference to their Use in Dyeing Processes, Bell, V A and Lewis, D M, Proceedings of the 5th International Wool Research Conference, Aachen, III (1975) 595-606
3. Advanced Technology for Bright, Fast Colours: Pad Batch Dyeing of Wool, Lewis, D M and Seltzer, I, *Text. J. of Australia*, **47** (1972) 24
4. A Combined Cold Dyeing and Resin Finishing Process for the Production of Washable Wool Fabrics, Lewis, D M, *Textilveredlung*, **8**, (1973) 151-158
5. Pad-Batch Dyeing of Wool Fabric - Development of an Industrial Process, Gibson, J D M, Lewis, D M and Seltzer I, *J.S.D.C.*, **86** (1970) 298-303
6. The Finishing of Wool Tops with Bunte Salt Polyamide Polymers, Cockett, K R F and Lewis, D M, *J.S.D.C.*, **92** (1976) 399-406
7. The Reaction of Reactive Dyes with Thiourea, Bell, V A and Lewis, D M, *J.S.D.C.*, **93** (1977) 140-141
8. The Dyeing of Wool Treated with Hercosett 57, Lewis, D M, and Seltzer, I, *Appl. Polym. Symp.*, **No. 18** (1971) 437-446
9. Dyeing Wool Piece Goods, Lewis, D M, *Text. Manuf.*, **95** (1969) 250
10. Print-Batch (Cold) - A New Process for Wool, Lewis, D M, and Seltzer, I, *J.S.D.C.*, **88** (1972) 327-329
11. Wool Dyeing, Lewis, D M, *Rev. of Prog. in Coloration*, **8** (1977) 10-23
12. The Reaction of -chloroacetyl-amino Dyes with Wool, Lewis, D M, Rattee, I D and Stevens, C B, *Proceedings of the 3rd International Wool Text. Res. Conf.*, Paris, CIRTEL III (1965) 305-317
13. Concentration Effects in Chromatographic Analysis, Lewis, D M, Rattee, I D and Stevens, C B, *Nature*, **208** (1965) 269-271
14. Wool Dyeing - A Path to Progress, Lewis, D M, and Seltzer, I, *J.S.D.C.*, **84** (1968) 101-607
15. Pad-Batch Dyeing of Wool with Reactive Dyes, Lewis, D M and Seltzer, I, *J.S.D.C.*, **84** (1968) 501- 507

16. The Dyeing of Wool with Premetallised Dyes, Lewis, D M, *Wool Science Review*, **54** (1977) 1-16.
17. Färbung von Wolle nach dem Foulardier - Kaltverweil Verfahren, Lewis, D M and Seltzer, I, *Textilveredlung*, **6** (1971) 87-92.
18. Dyeing Wool with Reactive Dyes, Lewis, D M, *Int. Dyer*, **142** (1969) 35
19. New Possibilities for the Low Temperature Dyeing of Wool, Graham, J F, Holt, R R D and Lewis, D M, *Proceedings of the 5th Int. Wool Textile Res. Conference, Aachen*, V (1975) 200-210
20. Production of Fast Dyeings on Resin Treated Wool, Lewis, D M and Seltzer, I, *J.S.D.C.*, **88** (1972) 93-100
21. Correspondence on above paper, Lewis, D M, *J.S.D.C.*, **88** (1972) 298
22. Development in Wool Dyeing with Particular Reference to Knits, Graham, J F and Lewis, D M, *Knitting Times*, **39** (1970) 22.
23. Combined Dyeing and Finishing of Wool, Lewis, D M, *International Dyer*, **156** (1976) 440
24. Continuous Dyeing and Polymer Shrink-Resist Processes for Wool, Lewis, D M, *J.S.D.C.*, **93** (1977) 105-113
25. The Development and Application of Reactive Dyes for Wool, Part I, Lewis, D M, *Wool Sci. Rev.*, **49** (1974) 13. Part II, Lewis, D M, *Wool Sci. Rev.*, **50** (1974) 22
26. The Pad-Batch (Cold) Dyeing Process for Wool, Gibson, J D M, Lewis, D M, and Seltzer, I, *Appl Polym. Symp.*, No. 18, (1971) 419-428
27. Shrink Resist Finishing of Wool Knits, Cockett, K R F. and Lewis, D M, *Text. Chem. and Colourist*, **10** (1978) 238-240
28. Transfer Printing of Wool, Bell, V A, Brady, P R, Cookson, P G and Lewis, D M, *Text. Chem. and Colorist*, **11** (1979) 100-105
29. Sublimation Transfer Printing of Surfactant Pretreated Wool, Bell, V A, Brady, P R, Cookson, P G, Fincher, K W, and Lewis, D M, *J.S.D.C.*, **97** (1981) 128
30. Sublimation Transfer Printing of Wool, Brady, P R, Cookson, P G, Fincher, K W, and Lewis, D M, *J.S.D.C.*, **96** (1980) 188-189
31. The Colouration of Wool and Wool Blends with Disperse Dyes, Bell, V A and Lewis, D M, *J.S.D.C.*, **94** (1978) 158-159

32. The Coloration of Wool with Disperse Dyes, Bell, V A and Lewis, D M, *J.S.D.C.*, **94** (1978) 507-513
33. The Development of Polymer Exhaust Aqueous Batch Processes to Meet the Superwash Standard, Cockett, K R F, Kettlewell, R, Lewis, D M, and Smith, P, *J.S.D.C.*, **96** (1980) 214-223
34. Applying Shrink Resist Polymers to Wool Fabrics at Room Temperature using Pad-Store Methods, Lewis, D M, *Text. Res. J.*, **52** (1982) 580-586
35. Developments in Wool Printing, Bell, V A, Lewis, D M, *Text. Res. J.*, **53** (1983) 125-141
36. The Dyeing of Wool with Reactive Dyes in the Presence of Trichloroacetic Acid, Lewis, D M, *J.S.D.C.*, **97** (1981) 365-369
37. The Dyeing of Wool with Reactive Dyes, Lewis, D M, *J.S.D.C.*, **98** (1982) 165-175
38. Sublimation Transfer Printing of Wool with Metallizable Dyes, Brady, P R Cookson, P G Fincher, K W and Lewis, D M, *J.S.D.C.*, **98** (1982) 398-403
39. Photo-acoustic Spectroscopy for the Study of the Adsorption of Dyes on Wool Fabrics, Davidson, R S, King, D, Duffield, P A, and Lewis, D M, *J.S.D.C.*, **99** (1983) 123-126
40. The Benzoylation of Wool and its Effect on Dye Affinity and Shrink Resistance, Lewis, D M, and Pailthorpe, M T, *J.S.D.C.*, **99** (1983) 354-357
41. The Modification of Wool with Reactive Hydrophobes, Lewis, D M, and Pailthorpe, M T, *J.S.D.C.*, **100** (1984) 56-63
42. Evaluation of a Series of Reactive Ionic, Nonionic, and Hydrophobic Compounds as Antistatic Agents for Wool, Lewis, D M, and Pailthorpe, M T, *Text. Res. J.*, **54** (1984) 279-288
43. Einige Aspekte des Drucks von Wolle und Wollmischungen, Bell, V A, Lewis, D M, Brady, P R, Cookson, P G, and Pailthorpe, M T, *Schriftenreihe des Deutschen Wollforschungsinstitutes*, **93** (1984) 102
44. Recent Developments in Wool Dyeing and Printing, Lewis, D M, and Pailthorpe, M T, *SDCANZ Conference Paper*, Feb. 1983 (Sydney)
45. An Investigation into New Pretreatments for the Printing of Wool and Wool-Blend Fabrics, Bell, V A, Brady, P R, Bryne, K M, Cookson, P G, Lewis, D M, and Pailthorpe, M T, *AATCC National Technical Conference Book of Papers* (1983) p. 289

46. Chemical Treatments Designed to Modify the Affinity of Wool for Dyes, Bell, V A, Lewis, D M, and Pailthorpe, M T, *J.S.D.C.*, **100** (1984) 223-231
47. Preparation and Characterisation of Dye-Metal Complexes formed during the Keratrans Process, Lewis, D M, Otterburn, M S and Savage, D A, *J.S.D.C.*, **100** (1984) 264-267
48. The Yellowing and Bleaching of Wool, Duffield, P A and Lewis, D M, *Rev. Progress in Colouration*, **15** (1985) 38-51
49. The Photostability of Fluorescent Whitening Agents on Wool, Davidson, R S, King, D, Lewis, D M, and Jones, S K R, *J.S.D.C.*, **101** (1985) 291
50. Reactive Dyes for Wool; Chemistry, Application and Market Status, *Arbeitstagung Lecture*, Aachen 1985
51. Practical Dye-Resist Treatments for Printing and Cross Dyeing of Wool, Bell, V A, Lewis, D M, and Pailthorpe, M. T, *Proceedings of the 7th Int. Wool Text. Res. Conf.*, Japan, 1985
52. Reactions Between Sulphamic Acid and Wool, Cameron, B A, Lewis, D M, and Pailthorpe, M T, *Proceedings of the 7th Int. Wool Text. Res. Conf.* Japan 1985
53. Pad-Batch Polymer Shrink-Resist Processes for Wool Fabrics, Bell, V A, Byrne, K M, Cookson, P G, Lewis, D M, and Rushforth, M A, *Proceedings of the 7th Int. Wool Text. Res. Conf.*, Japan 1985
54. Photoelectron Spectroscopy and the Surface Chemistry of Wool, Carr, C N, Ho, S F, Lewis, D M, Owen, E D and Roberts, M W, *J. Text. Inst.*, **76** (1985) 419
55. Reaktivfarbstoffe für Wolle: Chemie, Anwendung und Markt-Position, Lewis, D M, *Melliand Textilber*, **67** (1986) 717-723
56. The Photosensitizing Properties and Photostability of Stilbene Fluorescent Whitening Agents, Davidson, R S, Ismail, G and Lewis, D M, *J.S.D.C.*, **103** (1987) 261-264
57. Retardation of the Photoyellowing of Untreated Wool and Wool Treated with Fluorescent Whitening Agents by the Action of Reducing Agents, Davidson, R S, Ismail, G and Lewis, D M, *J.S.D.C.*, **103** (1987) 308-313
58. The Great Debate, Lewis, D M, *J.S.D.C.*, **103** (1987) 285
59. Insectproofing of Wool, Lewis, D M and Shaw, T, *Review of Progress in Colouration*, **17** (1987), 86-94
60. The Natural Fluorescence of Wool, Collins, S, Davidson, R S, Greaves, P H, Healey, M and Lewis, D M, *J.S.D.C.*, **104** (1988) 348-352

61. The Efficiency of Reaction between Reactive Fluorescent Whitening Agents and Wool, Davidson, R S, Ismail, G and Lewis, D M, *J.S.D.C.*, **103** (1987) 86-93
62. Dyeing for a Living, Lewis, D M, *Leeds University Review*, **31** (1988/89) 93-120.
63. Dyeing for a Living, Lewis, D M, *J.S.D.C.*, **105** (1989) 119-128
64. Analysis of Wool Lipids Using Thin-Layer Chromatography (TLC) with Flame Ionization Detection (FID), Leaver, I H, Lewis, D M and Westmorland, D J, *Textile Research Journal*, **58** (1988) 593-600
65. Improved Cellulose Dyeability by Chemical Modification of the Fibre, Lei, X P and Lewis, D M, *Proceedings of the AATCC Symposium*, Global Trends in Dyeing Science and Technology, (1989) 5-14
66. Testing the Washability of Woven Wool Fabrics, Grundell, P, Moreland, A, Lewis, D M, and Ren, J, *Wool Record*, April 1988 p.79
67. Some Aspects of the Photochemistry of Fibrous Proteins, Lewis, D M, *Colourage*, **36** (1989) 25-31
68. Damage in Wool Dyeing, Lewis, D M, *Review of Progress in Coloration*, **19** (1989) 49-56
69. The Effect of Reactive Dyes on Damage in Wool Dyeing, Lewis, D M, *Journal of Society of Dyers and Colourists*, **106** (1990) 270-274
70. Modification of Cotton to Improve its Dyeability, Part I - Pretreating Cotton with Reactive Polyamide-epichlorohydrin Resin, Burkinshaw, S M, Lei, X-P and Lewis, D M, *Journal of Society of Dyers and Colourists*, **105** (1989) 391-398
71. Reactive Dyes and Wool Damage, Lewis, D M, and Smith, S M, *Proceedings 8th International Wool Textile Research Conference*, **IV** (1990) 177-186
72. Dyeing and Wet Processing of Wool, Lewis, D M, *Proceedings 8th International Wool Textile Research Conference*, **IV** (1990) 1-49
73. The Preparation and Analysis of Disulphides as Model Compounds for Cystine in Wool, Douthwaite, F J , Lewis, D M, and Schumacher-Hammadat, U, *Proceedings 8th International Wool Textile Research Conference*, **I** (1990) 326-335
74. Improved Cellulose Dyeability by Chemical Modification of the Fibre, Lewis, D M, and Lei, X-P, *Textile Chemist and Colorist*, **21** (No. 10) (1989) 23-29
75. Lewis, D M and Lei, X-P, *Proceedings International Federation of Textile Chemists and Colourists, Lucerne Conference* (1990)

76. Modification of Cotton to Improve its Dyeability. Part 2 - Pretreating Cotton with a Thiourea Derivative of Polyamide-epichlorohydrin Resins, Burkinshaw, S M, Lei, X-P, Lewis, D M, Easton, J R, Parton, B and Phillips, D A S, *J.S.D.C.*, **106** (1990) 307-315
77. Modification of Cotton to improve its Dyeability. Part 3 - Polyamide-epichlorohydrin Resins and their Ethylene Diamine Reaction Products, Lei, X-P and Lewis, D M, *J.S.D.C.*, **106** (1990) 352-356
78. The Use of FTIR in the Study of Surface Chemistry of Hair Fibres, Joy, M and Lewis, D M, Proc. of the Society of Cosmetic Scientists Conference, Harrogate, 1990
79. New Methods for Improving the Dyeability of Cellulose Fibres with Reactive Dyes, Lewis, D M and Lei, X-P, *J.S.D.C.*, **107** (1991) 102-109.
80. The Role of Vinylsulphonyl Reactive Dyes in Prevention of Wool Damage, Lewis, D M and Smith, S M, *J.S.D.C.*, **107** (1991) 357-362
81. The Dyeing Behaviour of Cotton Modified with Chloropropionyl Chloride and Related Compounds, Lei, X-P and Lewis, D M, *Dyes and Pigments*, **16** (1991) 273-289
82. The Use of Fourier Transform Infra-red Spectroscopy in the Study of the Surface Chemistry of Hair Fibres, Joy, M and Lewis, D M, *International J. of Cosmetic Science*, **13** (1991) 249-261
83. Reactions of β -Sulphato-ethylsulphone Crosslinking Agent with Wool, Part 1: HPLC Analysis and Stripping Tests, Otterburn, M S, Smith B, Spedding, P, Lewis, D M and Allanach, D, *Text. Res. J.*, **61** (1991) 705-714
84. Some Aspects of the Photochemistry of Fibrous Proteins, in 'Leather - its Composition and Changes with Time', Lewis, D M, publ. The Leather Conservation Centre, (1991) 60-65
85. Analysis of Reactive Dyes and Related Derivatives using High-Performance Capillary Electrophoresis, Croft, S N and Lewis, D M, *Dyes and Pigments*, **18** (1992) 309-317
86. The Production of Level Fast Dyeings on Wool with Nucleophilic Dyes and a Crosslinking Agent, Lei, X-P., Lewis, D M and Wang, Y-N., *J.S.D.C.*, **108** (1992) 383-387
87. Method of Setting Hair, Lewis, D M, USP 5,071,641, Dec 10 (1991)
88. Neutral-fixing Reactive Dyes for Cotton. Part 1 - Synthesis and Application of Quaternised s-Triazinyl Reactive Dyes, Croft, S N, Lewis, D M, Orita, R and Sugimoto, T, *J.S.D.C.*, **108** (1992) 195-199

89. The Aftertreatment of Wool Dyeings to Obtain Improved Wet Fastness, Lewis, D.M., *J.S.D.C.*, **108** (1992) 317-324
90. Reactions of β -Sulphato-ethylsulphone Crosslinking Agent with Wool. Part II: Mechanism from Solubility and Swelling Tests, Smith, B M, Spedding, P L, Otterburn, M S and Lewis, D.M., *Textile Research Journal*, **62** (1992) 309-316
91. Ancillary Processes in Wool Dyeing, Lewis, D.M., Wool Dyeing: publ S.D.C. (1992) 111-136
92. Dyeing Wool with Reactive Dyes, Lewis, D.M., Wool Dyeing: publ S.D.C. (1992) 222-256
93. Environmental Aspects of Wool Dyeing, Lewis, D.M., New Wool Fabrics Conference, Huddersfield, July 1991
94. An FTIR Spectroscopic Study of the Photodegradation and Thermal Degradation of Wool, Carr, C M, and Lewis, D.M., *J.S.D.C.*, **109** (1993) 21-24
95. Novel Pigment Binders based on S-thiosulphate (Bunte Salt) Terminated Polyethers, Hinks, D, and Lewis, D.M., *J.S.D.C.*, **109** (1993) 106-111
96. Reactive Fibres: An alternative to Reactive Dyes, Lewis, D.M., and Lei, X-P, American Association of Textile Chemists and Colorists, Annual Conference & Exhibition, Atlanta, (1992): Book of Papers, 259-265
97. Reaction of Cystine Residues in Wool with Peroxy Compounds, Douthwaite, F J, Lewis, D.M., and Schumacher-Hamedat, U, *Text.Res.J.*, **63** (1993) 177-183
98. The Effect of Various Chromium Species on Wool Keratin, Lewis, D.M., and Yan, G, *J.S.D.C.*, **109** (1993) 193-198
99. Capillary Zone Electrophoresis in the Analysis of Dyes and Other Compounds Employed in the Dye Manufacturing and Using Industries, Burkinshaw, S M, Hinks, D, and Lewis, D.M., *J. of Chromatography*, **640** (1993) 413-417
100. New Possibilities to improve Cellulosic Fibre Dyeing Processes with Fibre-reactive Systems, Lewis, D.M., *J.S.D.C.*, **109** (1993) 357-364
101. Chrome Dyeing Using Cr(III) Salts, Lewis, D.M., and Yan, G, *J.S.D.C.*, **110** (1994) 281
102. The Natural Fluorescence of Wool, Collins, S, Davidson, R S, Hildenbach, M E C, and Lewis, D.M., *Dyes and Pigments*, **24** (1994) 151-169
103. The Use of Substantive Cross-linking Agents to Fix Nucleophilic Dyes on PolyamideFibres, Ho, Y-C and Lewis, D.M., American Association of Textile Chemists and Colorists, Annual Conference and Exhibition, Charlotte (1994) Book of Papers, 419-455

104. The Formation of Cysteine-S-sulphonate Groups in Wool and the Effect on Shrink-resistance, Douthwaite, F J, and Lewis, D M. *J.S.D.C.*, **110** (1994) 304-307
105. Improved Dyeing of Cellulosic Fibres, Broadbent, P A, Lei, X-P, and Lewis, D M, *Cellulosics: Materials for Selective Separations and Other Technologies*, (1993) 297-309, eds. Kennedy, J F, Phillips, G O, and Williams, P A, publ Ellis Horwood, ISBN 0-13-020298-3
106. Dyeability of Nylon Fibres, Lewis, D M, *Synthetic Fibre Materials*, (1994) 53-67, ed Brody, H., publ Longman, ISBN 0 582 06267 5
107. Oxidative Hair Dye Compositions and Processes Utilizing Leuco Vat Dyes, Lewis, D M, US Patent 5,364,415, Assignee Clairol Inc, New York, 15 Nov 1994
108. Reactions of β -sulphato Ethyl Sulphone Crosslinking Agent with Wool. Part III: Wool Fractionation and Amino Acid Analysis, Smith, B M, Spedding, P L, Otterburn, M S, and Lewis, D M, *Dyes and Pigments*, **26** (1994) 277-295
109. The Use of Fourier Transform Infrared (FTIR) Analysis to Study Chemical Changes in Wool Following a Sulphitolysis Treatment, Gomez, N , Erra, P, Lewis, D M, Willoughby, J H, *Book of Papers, ISF '94, Yokohama* (1994) 96
110. The Use of FT-IR for the Analysis of Reactive Dyes, Lewis, D M, and Gillingham, E L, *Book of Papers, ISF '94, Yokohama* (1994) 81
111. Dye-fibre Interactions, Lewis, D M, *Proceedings of the 12th International Farben Symposium, Bad Homburg*, 1994
112. Reactions of β -Sulphatoethyl Sulphone Crosslinking Agent with Wool. Part IV: Topochemistry of Reactions by Fractionation and Extraction, Smith, B M, Spedding, P L, Otterburn, M S and Lewis, D M, *Dyes and Pigments*, **28** (1995) 57-67
113. Reactions of β -Sulphatoethyl Sulphone Crosslinking Agent with Wool. Part V: Effects during Reactive Dyeing, Smith, B M, Spedding, P L, Otterburn, M S and Lewis, D M, *Dyes and Pigments*, **28** (1995) 101-122
114. Improved Fixation of Dyes on Polyamide Fibres. Part 1: Using 1,3,5-Triacryloylamino-Hexahydro-*s*-Triazine as a Crosslinking Agent, Ho, Y C and Lewis, D M, *Dyes and Pigments*, **28** (1995) 171-192
115. Level Fast Dyeing of Wool with Nucleophilic Aminoalkyl Dyes and Crosslinking Agents. Part 1: Using a Trifunctional Crosslinking Agent, Lei, X-P., Lewis, D M, and Wang, Y-N, *J.S.D.C.*, **111** (1995) 12-18
116. A New Approach to the Dyeing of Silk with Sulphatoethylsulphone Dyes, Lewis, D M and Shao, J Z, *J.S.D.C.*, **111** (1995) 146-150

117. The Use of FTIR to Investigate Modifications to Wool Treated with Sodium Sulphite and Cationic Protein Hydrolysate, Gomez, N, Julia, M R, Lewis, D M and Erra, P, *J.S.D.C.*, **111** (1995) 281-284
118. Improved Fixation of Dyes on Polyamide Fibres, Part 2: Use of the Reactive Crosslinking Agent, 2-Chloro-4,6 - Di(Aminobenzene-4-Sulphatoethylsulphone)-s-Triazine (XLC), Lewis, D M and Ho, Y C, *Dyes and Pigments*, **28** (1995) 237-259
119. Chrome Dyeing Using Anionic Trivalent Chromium Complexes as Mordants, Lewis, D M and Yan, G, *J.S.D.C.*, **111** (1995) 316-323
120. Dyeing in Non-aqueous Systems, Lewis, D M, Chapter 15 in Colour in Dyehouse Effluent, ed. P. Cooper, publ. Society of Dyers and Colourists, (1995) 195-197
121. Substituted Thiol Derivatives of Vinyl Sulphone Dyes, Lewis, D M and Smith, S M, *Dyes and Pigments*, **29** (1995) 275-294
122. Level Fast Dyeing of Wool with Nucleophilic Aminoalkyl Dyes and Crosslinking Agents, Lei, X-P, Lewis, D M and Wang, Y N, *J.S.D.C.*, **111** (1995) 385-390
123. The Effect of Environmental Legislation on Technological Innovation in the Colour-Making and Colour-Using Industries, Lewis, D M, Book of Papers, AATCC Intl. Dyeing Symposium (Lake Buena Vista), (1995) 99-114
124. Fixation of Aspartyl-triazine Dyes on Cellulosic Fibres, Lewis, D M and Gillingham, E L, American Association of Textile Chemists and Colorists, Annual Conference and Exhibition, Atlanta, (1995): Book of Papers, 536-544
125. Improved Fixation of Dyes on Polyamide Fibres, Part 3: Using 2-Chloro-4,6-di(aminobenzene-4-vinylsulphone)-s-triazine as an Aftertreatment of Nucleophilic-Aminoalkyl Dyes, Lewis, D M and Ho, Y C, *Dyes and Pigments*, **30** (1996) pp 301-314
126. Improved Fixation of Dyes on Polyamide Fibres, Part IV: The Use of a Nucleophilic Aminoethylsulphonyl Cationic Dye for Dyeing Nylon Followed by a Fixation Aftertreatment with 2-chloro-4,6-di(aminobenzene-4'-sulphatoethylsulphone)-s-triazine [XLC], Lewis, D M and Ho, Y C, *Dyes and Pigments*, **31** (1996) pp 111-129
127. Dyestuff-fibre Interactions, Lewis, D M, *International Journal of Cosmetic Science*, **18** (1996) pp 123-135
128. Fixation of Aspartyl-Triazine Dyes on Cellulosic Fibres, Lewis, D M and Gillingham, E.L., *Text. Chem. and Col.*, **28** (1996) pp 76-80

129. Chrome Dyeing of Wool with Cr^{III} Salts and Organic Acids, Lewis D M and Yan, G, Proceedings of the International Wool Textile Research Conference, Biella III (1996) pp 39-51
130. A Universal Dye for All Fibres - Are Disperse Dyes Capable of Fulfilling this Vision? Broadbent, P J and Lewis, D M, International Federation of Textile Chemists and Colorists 17th Conference, Vienna, Book of Papers, (1996) pp 107-111
131. Dyeing Polyamide and Cellulosic Fibres by Covalently Fixing Nucleophilic Dyes with Crosslinking Agents, Lewis, D M, Lei, X-P, Ho, Y C, Jabbar, A and Wang, Y N, Colorchem 96 Conference, Spindlerlov Myln, Czech Republic, Book of Papers, (1996) pp 32-35
132. Resin Coatings for Capillaries Giving a Net Positive Charge and Great Potential for Customised Modification of Surface Properties, Burt, H, Lewis, D M and Tapley, K N, *Journal of Chromatography A*, **739** (1996) 367-371
133. Coated Versus Fused Silica Capillaries for the Separation of Inorganic and Organic Cations by Capillary Zone Electrophoresis, Burt, H, Lewis, D M and Tapley, K N, *Journal of Chromatography A*, **736** (1996) 265-272
134. Crosslinking Nucleophilic Dyes on Wool, Lei, X P, Lewis, D M, Shen, X M and Wang, Y N, *Dyes and Pigments*, **30** (1996) 271-281
135. Dyeing Nylon 6,6 with Vinylsulphone Reactive Dyes, Lewis, D M and MacDougall, W C, American Association of Textile Chemists and Colorists, International Conference and Exhibition, (Nashville), Book of Papers (1996) 284-295
136. Crosslinking Nucleophilic Dyes on Wool, Lei, X P, Lewis, D M Shen, X M and Wang, Y N, Proceedings 9th International Wool Textile Research Conference, Biella, III (1996) 152-160
137. The Use of Fourier Transform Infrared (FTIR) Analysis to Study Chemical Changes in Wool Following a Sulphitolysis Treatment, Erra, P, Gomez, N, Julia, M R, Lewis, D M and Willoughby, J P, Proceedings 9th International Wool Textile Research Conference, Biella, II (1996) 387-393
138. Reactive Cellulose Fibres Rather than Reactive Dyes, Lewis, D M and Fan, Q C, The Chemistry and Processing of Wood and Plant Fibrous Materials, (1996) 221-226, eds Kennedy, J F, Phillips, G O and Williams, P A, Woodhead Publishing Co, Cambridge, UK
139. FTIR Analysis to Study Chemical Changes in Wool Following a Sulphitolysis Treatment, Erra, P, Gómez, N, Julià, M R, Lewis, D M and Willoughby, J H, *Textile Research J.*, **67** (1997) 397-400

140. FTIR Investigation of the Damage Produced on Human Hair by Weathering and Bleaching Processes: Implementation of Different Sampling Techniques and Data Processing, Signori, V and Lewis, D M, *International Journal of Cosmetic Science*, **19** (1997) 1-13
141. A Universal Dye for All Fibres - Are Disperse Dyes Capable of Fulfilling this Vision? Lewis, D M and Broadbent, P J, *Journal of the Society of Dyers and Colourists*, **113** (1997) 159-164
142. Solubility of Disperse Dyes in Supercritical Carbon Dioxide, Ozcan, A S, Clifford, A A, Bartle, K D and Lewis, D M, *J.Chem.Eng.Data*, **42(3)** (1997) 590-592
143. Coloration in 2000 AD, Lewis, D M, *Sen-i-Gakkaishi*, **53** (1997) 213-219
144. Surface Layer of Wool. I. Dityrosine Synthesis and Characterization, Stewart, K, Spedding, P L, Otterburn, M S and Lewis, D M, *Journal of Applied Polymer Science*, **66** (1997) 2359-2363
145. Surface Layer of Wool. II. Dityrosine in Wool, Stewart, K, Spedding, P L, Otterburn, M S and Lewis, D M, *Journal of Applied Polymer Science*, **66** (1997) 2365-2376
146. Dyeing of Cotton Fibres with Disperse Dyes in Supercritical Carbon Dioxide, Özcan, A S, Clifford, A A, Bartle, K D and Lewis, D M, *Dyes and Pigments*, **36(2)** (1998) 103-110
147. Durable Press Finishing of Cotton with Polycarboxylic Acids. 1. Preparation of Thiosuccinyl-s-triazine, Lewis, D M and Voncina, B, *Journal of Applied Polymer Science*, **66** (1997) 1465-1474
148. Durable Press Finishing of Cotton with Polycarboxylic Acids. II. Ester Crosslinking of Cotton with Dithiosuccinic Acid Derivative of s-Triazine, Lewis, D M and Voncina, B, *Journal of Applied Polymer Science*, **66** (1997) 171-177
149. Dyeing Nylon 6,6 with Vinylsulphone Reactive Dyes, Lewis, D M and MacDougall, W C, *Textile Chemist and Colorist*, **30** (1998) 31-35
150. The Use of Fourier Transform Infrared (FT-IR) Spectroscopy to Study the State of Heterobifunctional Reactive Dyes, Lewis, D M and Wang, J C, *Dyes and Pigments*, **39** (1998) 111-123
151. Reactive Fibres – An Alternative to Reactive Dyes, Lewis, D M and Lei, X-P, *Advances in Colour Science and Technology*, **1** (1998) pp 40-45
152. Investigating the Photo-oxidation of Wool Using FT-Raman and FT-IR Spectroscopies, Jones, D C, Carr, C M, Cooke, W D and Lewis, D M, *Textile Research Journal*, **68** (1998) 739-748

153. The Preparation of Bunte Salt-terminated Surface-active Agents and Their Use in Wool Fabric Shrinkproofing, Lewis, D M, *Journal of the Society of Dyers and Colourists*, **115** (1999) 59-61
154. The Use of a Crosslinking Agent to Achieve Covalent Fixation of Hydroxyethylsulphone Dyes on Cotton, Lewis, D M and Yao, J, *Advances in Colour Science and Technology*, **1** (1999) 93-98
155. Discharge Printing, Du, F. and Lewis, D M, *Advances in Colour Science and Technology*, **2** (1999) 104-108
156. The Application of Vat Pigment Dispersions and Leuco Vat Acid Dispersions to Polyester Fibres, Chevli, S N and Lewis, D M, *Advances in Colour Science and Technology*, **2** (1999) 148-151
157. Periodate Oxidized Cellulose for Dyeing with Alkylamino Dyes, Lewis, D M and Tapley, K N, *Textile Chemist and Colorist*, **31** (1999) 20-26
158. Modification of Cotton Cellulose with Sodium Benzoylthioglycollate and its Effect on its Dyeability with Disperse Dyes, Part 1: Synthesis and Characterisation of Sodium Benzoylthioglycollate, Broadbent, P J and Lewis, D M, *Dyes and Pigments*, **43** (1999) 51-58
159. Preparation of the Ethanolamine Derivative of C1 Reactive Red 3 and Its Covalent Fixation to Cotton by Crosslinking with a Polycarboxylic Acid, Lewis, D M and Voncina, B, *Colour Science '98*, **2** (1999) 231-235
160. Dyestuff-fibre Interactions, Lewis, D M, *Review of Progress in Coloration*, **28** (1998) 12-17
161. The Use of a Crosslinking Agent to Achieve Covalent Fixation of Hydroxyethylsulphone Dyes on Cotton, Lewis, D M and Yao, J, AATCC International Conference (Philadelphia), Book of Papers, (1998) 375-383
162. Recent Developments in Reactive Dye Chemistry and Processes for their Application, Lewis, D M, Proceedings of the International Symposium on Dyeing and Finishing of Textiles (Wakayama), (1998) 96-114
163. Water Pollution and Dyeing Processes, Lewis, D M, Proceedings of International Conference – Water and Textiles (University of Huddersfield), (1999) 91-105
164. Reactive Dyes and Dyeing: A Critical Review, English, R J and Lewis, D M, *Colour Science '98*, **2** (1999) 1-21
165. Polyalkylphosphonate Reactive Dyes for Cellulosic Fibres, Gillingham, E, Lewis, D M and Srikulkit, K, *Colour Science '98*, **2** (1999) 30-42
166. Coloration in the Next Century, Lewis, D M, *Review of Progress in Coloration*, **29** (1999) pp 23-28

167. Easy-care Finishing of Silk Fabric with a Polyphosphonic Acid Crosslinking Agent, Lewis D M and Shao J Z, *Advances in Colour Science and Technology*, **3**, (2000) pp 7-10
168. The Synthesis and Application of a New Reactive Dye Based on Disulphide-bis-ethylsulphone, Lewis D M, Renfrew A H and Siddique A A, *AATCC International Conference (Charlotte), Book of Papers*, (1999) pp 1-20
169. High Performance Capillary Zone Electrophoretic Analysis of a Thiosuccinyl-s-triazine Derivative, Voncina B, Lewis D M, *Advances in Colour Science and Technology*, **3**, (2000) pp 11-14
170. Chromium Distribution in Wool by Electron Microscopy and X-ray Energy Dispersive Analysis, Lewis D M, Yan G, Julia R M, Corderch L and Erra P, *Textile Research Journal*, **70**, (2000) pp 315-320
171. Dyeing Tosylated Cotton Fibres with Reactive Dyes and Dyes Containing Pendant Nucleophilic Groups, Arrowsmith L C and Lewis D M, *Advances in Colour Science and Technology*, **3**, (2000) pp 46-52
172. Cationic Fibre-reactive Dyes for Cellulosic and Polyamide Fibres, Hinks D, Burkinshaw S M and Lewis D M, *AATCC International Conference (Charlotte), Book of Papers*, (1999) pp 454-462
173. Reactive Dyeing Properties of Cotton Modified with 1-trimethylammonium-2-hydroxy-3-chloropropane chloride, Lewis D M and Wang H T, *Advances in Colour Science and Technology*, **3**, (2000) pp 60-64
174. Modification of Cotton Cellulose with Sodium Benzoylthioglycollate and its Effect on Dyeability with Disperse Dyes, Part 2: Application of Sodium Benzoylthioglycollate to Cotton Fabric, Broadbent P J and Lewis D M, *Dyes and Pigments*, **45**, (2000) pp 35-41
175. Covalent Fixation of Hydroxyethyl Sulphone Dye on Cotton by the Use of Crosslinking Agent Via a Pad-batch Process, Yao J and Lewis D M, *Journal of the Society of Dyers and Colourists*, **116**, (2000) pp 198-203
176. Dyeing of Modified Cotton Fibres with Disperse Dyes from Supercritical Carbon Dioxide, Özcan A S, Clifford A A, Broadbent P J and Lewis D M, *Journal of the Society of Dyers and Colourists*, **114**, (1998) pp 169-173
177. Crosslinking a Hydroxyethylsulphone Dye on Cotton to Achieve Covalent Fixation in a Long-liquor Process, Lewis D M and Yao J, *Journal of the Society of Dyers and Colourists*, **116**, (2000) pp 285-290
178. An FTIR Study of Anhydride Formation on Heating Butane-tetra-carboxylic Acid in the Presence of Various Catalysts, Gillingham E, Lewis D M and Voncina B, *Textile Research Journal*, **69**, (1999) pp 949-955

179. Recent Developments in Reactive Dyes: Chemistry and Application Processes, Lewis D M, *Advances in Colour Science and Technology*, **4**, (2001) pp 41-48
180. Synthesis of Hetero-bifunctional Reactive Dyes Containing a Disulphide-bis-ethylsulphone Cross-link and their Application to Cotton, Lewis D M and Siddique A A, *Advances in Colour Science and Technology*, **4**, (2001) pp 89-96
181. Analysis of Organic Cations by Capillary Electrophoresis, Lewis D M and Wang H T, *Advances in Colour Science and Technology*, **3**, (2000) pp 90-93
182. FTIR and FT-Raman Spectroscopic Studies of 2,4,6-trichloro-s-triazine and its Hydroxy Derivatives, Shen X M, Lewis D M and Tapley K N, *Advances in Colour Science and Technology*, **4**, (2001) pp 59-67
183. The Effect of Crosslinking Agents and Reactive Dyes in the Fibrillation of Lyocell, Karypidis M, Wilding M A, Carr C M and Lewis D M, *AATCC Review*, **1**, (2001) pp 40-44
184. The Characterisation of Cellulose Aldehyde Using Fourier Transform Infrared Spectroscopy, Fan Q G, Lewis D M and Tapley, *Journal of Applied Science*, **82**, (2001) pp 1195-1202
185. The Creativity Process – Luck or Good Judgement, Lewis D M, *Journal of the International Creativity and Communications Association (Japan)*, **6**, (2001) pp 1-6
186. Co-Application of Hydroxyalkyl Dyes and Polyphosphonic Acids to Cotton to Achieve Dye-Fibre Covalent Bonding, Gillingham E L, Lewis D M and Srikulkit K, *Coloration Technology (Society of Dyers and Colourists)*, **6**, (2001) pp 318-322
187. The Effect of Various Anti-setting Agents in Wool Dyeing. Part 1: Hydrogen Peroxide Based Systems, *Coloration Technology*, **118**, (2002) pp121-124
188. Reactive Dyeing Systems for Wool Fibres Based on Hetero-Bifunctional Reactive Dyes. Part 1: Application of Commercial Reactive Dyes, Cho H J and Lewis D M, *Coloration Technology (Society of Dyers and Colourists)*, **118**, (2002) pp 198-204
189. The Effect of Various Anti-Setting Systems in Wool Dyeing. Part 2: Sodium Maleate Based Systems, Kim J and Lewis D M, *Coloration Technology (Society of Dyers and Colourists)*, **118** (2002) pp 181-184
190. Chemical Modification of Cotton to Improve Fibre Dyeability, Wang H T and Lewis D M, *Coloration Technology (Society of Dyers and Colourists)*, **118**, (2002) pp 159-168
191. Chrome Dyeing of Wool Using Anionic Chromium (III) Complexes, Lewis D M, Hornett C, Smith P and Yan G, *Advances in Coloration Science and Technology*, **5**, (2002) 13-17

192. Wool Coloration – New Aspects and Opportunities, D M Lewis, *Proceedings of the World Wool Conference (Bradford)*, (2003) pp 103-123
193. Reactive Dyeing Systems for Wool Fibres Based on Hetero-Bifunctional Reactive Dyes. Part 2: Investigation of Dyeing Properties During the Dyeing Cycle, H J Cho and D M Lewis, *Coloration Technology*, **118** (2002) pp 220-225
194. Reactive Dyeing Systems for Wool Fibres Based on Hetero-Bifunctional Reactive Dyes. Part 3: Application of Specially Prepared Hetero-Bifunctional and Related Model Dyes on Wool, H J Cho and D M Lewis, *Coloration Technology*, **119** (2003) pp 1-9
195. Dyeing Natural Fibres in Supercritical Carbon Dioxide Using a Nonionic Surfactant Reverse Micellar System, K Sawada, T Takagi, M Ueda and D M Lewis, *Coloration Technology*, **118** (2002) pp 233-237
196. The Effect of Various Anti-Setting Systems in Wool Dyeing. Part 3: Sodium Thiocyanate Based Systems, J Kim and D M Lewis, *Coloration Technology*, **119** (2003) pp 108-111
197. The Effect of Various Anti-Setting Systems in Wool Dyeing. Part 4: Sodium Tetrathionate Based Systems, J Kim and D M Lewis, *Coloration Technology*, **119** (2003) pp 112-115
198. Modification of Cellulosic Fibres with Reactive Quaternary Ammonium Containing Cations; Evaluation by Capillary Zone Electrophoresis and Kjeldahl Nitrogen Determination, P Grooby, D M Lewis and M Clark, *Advances in Colour Science and Technology*, **6** (2003) pp 39-46
199. Thiocyanate Based Systems, J Kim and D M Lewis, *Coloration Technology*, **119** (2003) pp 108-111
200. Tetrathionate Based Systems, J Kim and D M Lewis, *Coloration Technology*, **119** (2003) pp 112-115
201. Quaternary Reactive Dyes Containing a Thioether-ethylsulphone Group, Part 1: Synthesis of a Trimethylammonium Ethylsulphide-ethylsulphone Dye, D M Lewis and L J Sun, *Coloration Technology*, **119** (2003) pp 286-291
202. Quaternary Reactive Dyes Containing a Thioether-ethylsulphone Group, Part 2: Application Methods for the Dyeing of Nylon, D M Lewis and L J Sun, *Coloration Technology*, **119** (2003) pp 327-330
203. Synthesis of Triethanolamine Trisulphate and its Application to Cellulosic Fabrics, D M Lewis and X F Zhao, *Coloration Technology*, **120** (2004) pp 172-179

204. Reactive Dyes, D M Lewis, *Kirk-Othmer Encyclopedia of Chemical Technology* – publ J Wiley & Sons Inc, (2004) pp 1-26
205. Nylon Dyeing, D M Lewis and D G Marfell, *Synthetic Fibre Dyeing* (publ Soc Dyers and Col), (2004) pp 82-121
206. Micellar Electrokinetic Capillary Chromatography Analysis of Phthalic Acid Isomers, M L A Rigout, D M Lewis and P J Broadbent, *J. Cap. Electrophoresis and Microchip Tech.*, **9**, (2005) pp 57-64
207. Modification of a Vinylsulphone Reactive Dye Using Diethanolamine-O-disulphate, D M Lewis and X F Zhao, *Coloration Technology*, **121**, (2005) pp 121-126
208. Ink-Jet Printing of Conductive Patterns on Textile Fabrics, Bidoki S M, McGorman D, Lewis D M, Clark M, Horler G, *Proceedings of AATCC International Conference and Exhibition*, (2004) pp 260-274
209. The Use of 1,3,5-triacroylamino-hexahydro-s-triazine (FAP) as a Cross-linker to Fix Dyes on Cellulosic Fibres, Lewis D M, and Jabbar A, *Advances in Colour Science and Technology*, **7**, (2005) pp101-107.
210. Synthesis of Reactive Dyes Based on the Bis-(N-carboxymethylamino)mono Quaternary-triazine-bis-ethylsulphone Reactive Group. Part 1: Application to Cotton Cellulose, Lewis D M and Siddique A A, *Coloration Technology*, **122**, (2006) pp 217-226
211. Use of Fourier Transform Infrared Spectroscopy to Follow the Heterocumulene Aided Thermal Dehydration of Phthalic and Naphthalic Acids, Rigout M L A and Lewis D M, *Applied Spectroscopy*, **60**, (2006) pp 1405-1413
212. Ink-jet Fabrication of Electronic Components, Bidoki S M, Lewis D M, Clark M, Vakarov A, Milner P A and McGorman D, *J. Micromech. Microeng.*, **17**, (2007) pp967-974
213. Improved Reactive Dyeing of Wool with Novel Trifunctional Reactive Dyes, Cho H J, Lewis D M and Jia B H, *Coloration Technology*, **123**, (2007) pp 86-95
214. Triazinylamino-alkylphosphonate Reactive Dyes for Cellulosic Fibres, Gillingham E L, Lewis D M, Nabi A and Srikulkit S, *Coloration Technology*, **123**, (2007) pp 178-183
215. The Use of a Novel Water-soluble Arylating Agent to Dye Wool and Wool Blends with Disperse Dyes, Broadbent P J, Clark M, Croft S N and Lewis D M, *Coloration Technology*, **123**, (2007) pp

PUBLISHED PATENTS

Cold Dyeing	GB 1,350,589
Novel Bunte Salts	GB 1,423,342

Bunte Salt Polymers	GB 1,423,341
Polytetrahydrofuran Resins	GB 1,480,213
Bunte Salt/Thiol Adducts	GB 1,587,406
Solvent Polymer Treatment	GB 1,593,858
Divalent Salt/Cloud Point For PPO Polymer Exhaustion	GB 2,005,322
Thiomaleate Ester Polymers	GB 1,565,353
New Polymer Exhaustion Method	GB 1,571,188
Trichloroacetic Acid in Dyeing	GB 2,028,382B
Transfer Printing	GB 2,025,471
Reactive Arylphobes	USA 4,563,189
Method of Setting Hair	USA 5,071,641 (1991)
Oxidative Hair Dyeing Compositions	USA 5,364,415 (1994)
Disperse Dye Pretreatment	GB 94/02403 (1994)
New Reactive Textile Dyes	WO 99/51682
New Reactive Textile Dyes	WO 99/51683
New Reactive Textile Dyes	WO 99/51684
New Reactive Textile Dyes	WO 99/51685
New Reactive Textile Dyes	WO 99/51686
New Reactive Textile Dyes	WO 99/51687
New Reactive Textile Dyes	WO 99/51688
New Reactive Textile Dyes	WO 99/51689
Fibre Reactive Arylating Agents	PCT/GB98/00526
Process for Improving Characteristics of Polyamide Fibres	US6258928B1 (2001)
Improvements in Reactive Dyes	WO 99/51682 (1999)

Improvements in Reactive Dyes	WO 99/51684 (1999)
Improvements in Reactive Dyes	WO 99/51686 (1999)
Improvements in Reactive Dyes	WO 99/51687 (1999)
Cotton Fabric with Durable Properties	WO 01/23660 (2001), US0026512
Vat Acid Dyeing of Textile Fibres	US6780205 (2004)
Dyestuffs (Customs & Excise)	GB 0406770.8 (2004)
Improvements in and Relating to Wool Treatment (Dec 05) Applicn. 0524371.2	
Improvements in and Relating to Image Articles (Dec 05) GB2414961	
Improvements in and Relating to Viologens	
The Use of Percarbamic Acid and Diacyl Peroxides and Precursors Therefor PCT/GB01/03656	
Rare Earth IR Absorbers (May 07) PCT/GB2007/001753	
Improvements in and Relating to Image Articles (Oct 2006) PCT/GB2006/004523	